

ACADEMIC
PRESSAvailable online at www.sciencedirect.com

SCIENCE @ DIRECT®

Journal of Combinatorial Theory, Series B 88 (2003) 17–27

Journal of
Combinatorial
Theory

Series B

<http://www.elsevier.com/locate/jctb>

A sufficient condition for planar graphs to be 3-colorable

O.V. Borodin¹ and A. Raspaud*LaBRI, Université Bordeaux I, 33405 Talence Cedex, France*

Received 26 June 2000

Abstract

Planar graphs without 3-cycles at distance less than 4 and without 5-cycles are proved to be 3-colorable. We conjecture that, moreover, each plane graph with neither 5-cycles nor intersecting 3-cycles is 3-colorable. In this conjecture, none of the two assumptions can be dropped because there exist planar 4-chromatic graphs without 5-cycles, as well as planar 4-chromatic graphs without intersecting triangles.

© 2002 Elsevier Science (USA). All rights reserved.

1. Introduction

As proved by Garey et al. [1], the problem of deciding whether a planar graph is 3-colorable is NP-complete. Therefore, it seems wise to discuss only sufficient conditions for a planar graph to be 3-colorable. As early as 1969, Havel [3] asked if there existed a constant C such that each planar graph with the minimal distance between triangles at least C was 3-colorable. In 1976, Steinberg conjectured that each planar graph without 4- and 5-cycles was 3-colorable (see Problem 2.9 [4,5]). Both problems remain widely open. One of the purposes of this paper is to pose the following.

Conjecture 1.1. *Every planar graph without intersecting 3-cycles and without 5-cycles is 3-colorable.*

E-mail addresses: brdnoleg@math.nsc.ru (O.V. Borodin), raspaud@labri.fr (A. Raspaud).

¹On leave of absence from the Institute of Mathematics, Siberian Branch, Russian Academy of Sciences, Novosibirsk 630090, Russia. A part of this research was supported by the Grants INTAS 97-1001 and 97-01-01075 of the Russian Foundation for Basic Research.

This [Bordeaux 3-color] conjecture if true is best possible in the sense that there exist 4-chromatic planar graphs without intersecting triangles (as shown by Havel [3], see Fig. 1) and also those without 5-cycles (see Fig. 2).

We also pose the following [strong Bordeaux] conjecture, which implies both Conjecture 1.1 and Steinberg’s one:

Conjecture 1.2. *Every planar graph without adjacent 3-cycles and without 5-cycles is 3-colorable.*

(By intersecting (adjacent) triangles we mean those with a vertex (an edge) in common.) Our second purpose is to make the first step in the direction of Conjectures 1.1 and 1.2.

Theorem 1.3. *Every planar graph with neither 3-cycles at distance less than 4 nor 5-cycles is 3-colorable.*

Informally speaking, this result shows that the “intersection” of Havel’s and Steinberg’s problems has a positive solution, i.e., each graph satisfying both Havel’s and Steinberg’s assumptions is 3-colorable. It also says that the main difficulty in Havel’s problem is hidden in the 5-cycles. Finally, it is the first positive result on 3-coloring that involves the minimal distance between triangles.

It was easier for us to prove the following fact:

Theorem 1.4. *Every proper 3-coloring of a face of size 3 or 7 in a connected plane graph that has neither a pair of 3-cycles at distance less than 4 nor a 5-cycle can be extended to a proper 3-coloring of the whole graph.*




Fig. 1. A 4-chromatic graph without intersecting triangles.




Fig. 2. A 4-chromatic graph without 5-cycles.




Fig. 3. Not extendable 3-coloring.

Fig. 3 represents two graphs such that the 3-coloring of the outside 6-face cannot be extended to the whole graph.

To deduce Theorem 1.3 from Theorem 1.4, suppose that G is a counterexample to Theorem 1.3 on the fewest vertices. Clearly, G is connected and by Grötzsch's theorem [2] has a 3-cycle, which is clearly a 3-face, f , by the minimality of G . By Theorem 1.4, a 3-coloring of f can be extended to a 3-coloring of G .

This formulation of Theorem 1.4 is not the strongest possible. As observed by a referee, (i) putting at least two vertices of degree 2 on an edge of the outside faces in Fig. 3 gives counterexamples with precolored faces of arbitrary size greater than 7 and, on the other hand, (ii) the problem of 3-color extension from a 4-face is reduced to that from a 7-face by putting three vertices of degree 2 on a boundary edge of the 4-face.

We denote the degree of a vertex v by $d(v)$ and the size of a face f by $r(f)$ (bridges are counted twice); a k -vertex is that of degree k , a $\geq k$ -vertex has the degree at least k , etc.

2. Proof of Theorem 1.4

2.1. Properties of a minimal counterexample

Let T be the minimum number of 3-cycles in the counterexamples to Theorem 1.4, and let N be the minimum number of vertices in the counterexamples having T cycles of length 3. Finally, let G have the fewest edges among the counterexamples to Theorem 1.4 having T 3-cycles and N vertices.

Suppose that a proper 3-coloring φ of a face f_0 in G cannot be extended to a proper 3-coloring of G . W.l.o.g., suppose that f_0 is the outside face. Denote the boundary of f_0 by $D = d_1 d_2 \dots d_{|D|}$.

(1) f_0 is a 7-face. Otherwise, we put four vertices of degree 2 on a boundary edge of the 3-face f_0 and extend φ to the new vertices. The graph G' obtained has fewer

3-cycles, is connected, and has neither 3-cycles at distance less than 4 nor 5-cycles. It follows, φ can be extended to G' , which gives also a proper 3-coloring of G .

(2) G is 2-connected; in particular, G has no 1-vertices. First suppose that the cycle D is not simple and d_1 is its cut-vertex. Then D consists of a 3-cycle $d_1d_6d_7$ and either (i) a 4-cycle $d_1d_2d_3d_4$, or (ii) two pendant edges (in two ways), or else (iii) a pendant 2-path. We put four 2-vertices on the edge d_1d_7 to obtain a 7-cycle C'_3 . In case (i), we put three 2-vertices on the edge d_1d_2 to obtain a 7-cycle C'_4 . Then we extend φ to C'_3 and C'_4 . Now we can extend φ to the interiors of C'_3 and C'_4 as each of them has fewer 3-cycles than G . This completes the proof in case (i); cases (ii) and (iii) are even easier.

Thus D is a simple cycle, and it is contained in a block of G . Hence, there is a pendant block B of G , with a cut-vertex v , such that $(B - v) \cap D = \emptyset$. The graph G' obtained from G by deleting $B - v$ has a 3-coloring that extends φ . In turn, B is 3-colorable, either by the Grötzsch theorem, or by the minimality of G , and we can choose a color for v in B the same as in G' . These two 3-colorings give a suitable 3-coloring of G .

(3) Each 2-vertex in G belongs to D .

(4) D is a simple 7-cycle without chords. Indeed, if D has a chord then, due to the absence of 5-cycles in G , this chord splits D into a 3-cycle and a 6-cycle. We delete the chord and extend φ to the remaining graph by the edge-minimality of G . The 3-coloring obtained is proper also for G .

(5) G has no separating 3-cycles and 7-cycles. If such a separating cycle S exists, we first extend φ from D to $G - \text{Int}(S)$, and then extend the obtained 3-coloring of the outside face S of $G - \text{Ext}(S)$ to the whole $G - \text{Ext}(S)$ (either by the minimality of G or by the Grötzsch theorem).

The following properties (5+) of G are easy consequences of (1–5) and will be used throughout the proof without reference.

- (5+) (a) Every 3-cycle is actually a 3-face,
 (b) each simple 7-cycle is either a 7-face or consists of a 3-face adjacent to a 6-face,
 (c) no 3-face is adjacent to a 4-cycle, and
 (d) no 3-face can share two edges with another face.
 (6) Each 4-face has a vertex in common with D . Suppose $f = wxyz$ is a face inside D . First observe that identifying x with z (or y with w) within f does not create a 3-cycle: this follows from the absence of 5-cycles in G .

Now suppose y is a vertex of f whose distance, $\rho(y)$, from the triangles of G is the least as compared to w, x and z , and let T be a triangle at distance $\rho(y)$ from y . Recall that f cannot share an edge with a 3-cycle. If $\rho(y)$ is at most 1, then the distance from x and z to T is $> \rho(y)$ due to the absence of 5-cycles in G (see Figs. 4 and 5). Hence




Fig. 4. $\rho(y) = 0 \Rightarrow z' \neq z$.




Fig. 5. $\rho(y) = 1 \Rightarrow zz' \notin E$.

$\rho(y) = 1$ implies $\rho(x) \geq 2$ and $\rho(z) \geq 2$, and if $\rho(y) = 0$ then each of x, z lies at distance 1 from T and at distance at least 3 from each other 3-face of G .

It follows that in both cases identifying x with z does not create a pair of triangles at distance less than 4.

We next show that identifying x with z does not create a 5-cycle $xv_1 \dots v_4$. Suppose the contrary, and by symmetry between y and w (the minimality of y is not used anymore) we can assume w.l.o.g. that y lies non-strictly inside the 7-cycle $zw xv_1 \dots v_4$. Due to (5), y must coincide with one of v_1, \dots, v_4 , but this is impossible due to the absence of 5-cycles.

Hence, identifying x with z cannot create a 5-cycle or a pair of 3-cycles at distance less than 4. Also observe that since neither x nor z belongs to D by assumption, it follows that the coloring φ of D remains proper, because there are only two ways to spoil φ by contraction: (i) to identify two vertices of D colored differently, or (ii) to join by an edge two vertices of D colored the same.

So, identifying x with z yields a graph G^* which has a 3-coloring that extends φ , and this coloring trivially gives a 3-coloring of G that extends φ , a contradiction.

(7) G cannot have a 4-face sharing two consecutive edges with D . Suppose $f = wxyz$ is a 4-face such that $\{x, y, z\} \in D$. Then $w \notin D$ because of the absence of 5-cycles in G . We extend φ to w by putting $\varphi(w) = \varphi(y)$. Clearly, φ is still proper since w cannot be adjacent to a vertex from $D - \{x, z\}$ due to the absence of 5-cycles. So, φ can be extended from the outside face of $G - y$ to the whole $G - y$, which immediately gives a 3-coloring of G ; a contradiction.

(8) G cannot have a 4-face $f = wxyz$ sharing precisely one edge, xy , with D . Suppose that $x = d_3$ and $y = d_2$. First observe that $z \notin D$: Due to (7), $z \neq d_1$, and if $z = d_i$ then we have a 5-cycle whenever $5 \leq i \leq 7$ and a separating 3-cycle for $i = 4$. By symmetry, $w \notin D$.

Due to the same argument as in proving (6), we can assume that identifying x with z does not create a 5-cycle nor a pair of 3-cycles at distance less than 4. To obtain a contradiction, it remains to show that this operation leaves the coloring φ on D proper.

Since z is inside D , we should only exclude the possibility to make φ improper because of the edge zd_t . By (5+), $1 \neq t \neq 3$. Suppose $t = 7$. Then we have a 7-cycle $d_4 \dots d_7 z y x$, which cannot be separating due to (5). It follows that $w \in D$, a contradiction. Clearly, $6 \neq t \neq 5$ because of the absence of 5-cycles. Finally, if $t = 4$, then identifying z with x does not actually create new adjacencies due to the presence of the edge xd_4 (we simply delete the edge zd_4).

Thus, we can extend φ to the graph with x and z identified, and hence to G itself; a contradiction.

(9) If a 3-vertex y is incident with a 6-face $f = xyzw_1w_2w_3$ and a 3-face $f' = xyt$, where $\{x, y, z\} \cap D = \emptyset$, then:

- (a) $d(t) > 3$, and
- (b) either $d(x) > 3$ or $t \in D$.

We first prove that

- (c) the third face $f'' = tyz \dots$ incident with y has size 6.

To this end, let us observe that deleting y followed by identifying x with z does not create a 3-cycle due to the absence of 5-cycles in G . Next, observe that due to the presence of the 3-face f' , this operation does not create two 3-cycles at distance less than 4. Since both x and z are inside D , their identification does not affect the coloring φ of D . Thus, the only obstacle for identifying x with z is creating a 5-cycle $xv_4 \dots v_1$. Suppose this has happened and consider the two cases.

Case 1: The path $w_1w_2w_3$ lies non-strictly inside the 7-cycle $xyzv_1 \dots v_4$. Due to (5), all w_i 's, where $1 \leq i \leq 3$, must coincide with some of v_j 's, where $1 \leq j \leq 4$. Clearly, $w_1 \neq v_2$ because of the absence of two 3-faces at distance at most 1. If w_1 coincides with v_3 or v_4 , we have a 5-cycle. Hence, $w_1 = v_1$, and, by symmetry, $w_3 = v_4$. Then w_2 coincides with v_2 or v_3 , and we have a 3-cycle too close to f' ; a contradiction.

Case 2: t lies non-strictly inside the 7-cycle $xyzv_1 \dots v_4$.

Similarly, t must coincide with one of v_1, \dots, v_4 . However, $t \neq v_4$ readily implies the presence of two adjacent 3-faces or a 5-cycle in G .

This completes the proof of (c).

Now we have a symmetric configuration shown in Fig. 6. To prove (a), suppose $d(t) = 3$ (by (2–4), $d(t) \geq 3$). Then t cannot lie in D , for otherwise x would be in D .




Fig. 6. Equation (9).

As follows from proving (9c), the only non-separating 7-cycle going through x and z goes also through t . We thus delete t along with y and destroy this last obstacle for identifying x with z .

So, identifying x with z followed by deleting y and possibly t , yields a graph which has a 3-coloring that extends φ , and this coloring trivially gives a 3-coloring of G that extends φ : if t was deleted, we first color t and then y ; otherwise we just color y .

This completes the proof of (a).

To prove (b), suppose that $t \notin D$. Then we apply (9) symmetrically, i.e., to the face f'' , which is a 6-face as proved in (c), rather than to f . (Now the roles of t and x are interchanged.) It follows from the symmetric version of (a) that $d(x) > 3$.

2.2. No minimal counterexample can exist

The rest of our proof consists in proving that the structural properties (1–9) of G are mutually incompatible. Euler's formula $|V(G)| - |E(G)| + |F(G)| = 2$ for G may be rewritten as

$$\sum_{v \in V(G)} (d(v) - 4) + \sum_{f \in F(G)} (r(f) - 4) = -8.$$

We set the *initial charge* of every vertex v of G to be $ch(v) = d(v) - 4$, that of every face $f \neq f_0$ to be $ch(f) = r(f) - 4$, and also set $ch(f_0) = r(f_0) + 4$. Clearly,

$$\sum_{x \in V(G) \cup F(G)} ch(x) = 0.$$

Now we use the discharging procedure, leading to the *final charge* ch^* , defined by applying the rules R0–R5 below. Since this procedure preserves the total charge, we have

$$\sum_{x \in V(G) \cup F(G)} ch^*(x) = 0.$$

The rest of our proof will consist in showing that $ch^*(x) \geq 0$ whenever $x \in V(G) \cup F(G)$, and $ch^*(f_0) > 0$, with the obvious final contradiction ($0 > 0$).

R0. Each 3-face $f = xyz$ receives $\frac{1}{3}$ from each adjacent face, unless $d(x) = 3$, $d(y) \geq 4$, and $d(z) \geq 4$, in which case f receives from adjacent faces $\frac{1}{6}$ across each of the edges xy , xz and $\frac{2}{3}$ across yz .

R1. Every 3-vertex $v \notin D$ receives $\frac{1}{3}$ from each incident face, unless v is incident with one 3-face, in which case v receives $\frac{1}{2}$ from each of the two >3 -faces.


R2. Every 2-vertex receives $\frac{5}{3}$ from f_0 and $\frac{1}{3}$ from the other (internal) incident face.

R3. f_0 gives 1 to every incident vertex of degree at least 3.


R4. Every vertex $d_2 \in D$ with $d(d_2) \geq 4$ gives 1 to each incident face not incident with the edges d_1d_2 and d_2d_3 . Furthermore, if 1 is given to a 3-face $f = d_2xy$ such that $x \notin D$, $y \notin D$, then this 1 is transferred by f across the edge xy to the neighbor face.

R5. Each ≥ 8 -face $f \neq f_0$ gives $(r(f) - 8)/2$ to f_0 .


R0.




R1.




R2.




R3.



R4.



R5.



(10) If $v \in V(G)$ then $ch^*(v) \geq 0$.

First suppose $v \notin D$. If $d(v) = 3$, then by R1 $ch^*(v) = 3 - 4 + 3 \times \frac{1}{3} = 0$ when v is not incident with a 3-face and $ch^*(v) = 3 - 4 + 2 \times \frac{1}{2} = 0$ otherwise. If $d(v) \geq 4$, then $ch^*(v) = ch(v) = d(v) - 4 \geq 0$.

Now suppose $v \in D$. If $d(v) = 2$ then $ch^*(v) = 2 - 4 + \frac{1}{3} + \frac{5}{3} = 0$ by R2. If $d(v) \geq 3$, then v receives 1 from f_0 by R3 and sends away $d(v) - 3$ units of charge by R4, so that $ch^*(v) = d(v) - 4 + 1 - 1 \times (d(v) - 3) = 0$.

(11) If $f \in F(G)$ and $f \neq f_0$, then $ch^*(f) \geq 0$.

If $r(f) = 3$, then its initial charge is -1 and f receives either $\frac{1}{3} + \frac{1}{3} + \frac{1}{3}$ or $\frac{2}{3} + \frac{1}{6} + \frac{1}{6}$ by R0, so that $ch^*(f) \geq 0$. (If f appears in R4, then f can only win from that in total.)

If $f = wxyz$, i.e., $r(f) = 4$, then its initial charge is 0. By (6), f has a vertex, say y , in common with D . Then neither x nor z belongs to D due to (8) and (4). It follows that $d(y) \geq 4$ and f receives 1 from y by R4. After giving $\frac{1}{3}$ by R1 to at most three incident vertices, f has $ch^*(f) \geq 0$. (By the absence of 5-cycles, f cannot give $\frac{1}{2}$ according to R1 or give anything to a neighbor 3-face by R0.)

We now consider the most difficult case $r(f) = 6$. Suppose $f = v_1 \dots v_6$ and recall that the initial charge of f is 2. Also recall that $f \neq f_0$ does not give anything to vertices of degree greater than 2 from D . If our f is not adjacent to a 3-face, then f gives at most $\frac{1}{3}$ to at most six vertices by R1, R2 so that $ch^*(f) \geq 2 - \frac{1}{3} \times 6 = 0$.

So suppose there is a 3-face $f' = tv_1v_2$. Such a face is unique, and it has only one edge v_1v_2 in common with f . If each of v_1 and v_2 has degree greater than 3 or belongs to D (and thus has degree at least 3, being incident with a 3-face), then f only gives $\frac{2}{3}$ across v_1v_2 by R0 and at most $\frac{1}{3}$ to at most four vertices by R1, and we are done.

We can thus assume that $d(v_2) = 3$ and $v_2 \notin D$. Now f sends at most $\frac{1}{3}$ across v_1v_2 and gives $\frac{1}{2}$ to v_2 . The total donation of f in the worst case cannot exceed $\frac{8}{3} : \frac{1}{2}$ to v_1 and $\frac{1}{3}$ to each of the four other incident vertices.

Let us first prove $ch^*(f) \geq 0$ assuming that at least one of the vertices v_1, v_3 belongs to D . Suppose $v_1 \in D$ (our argument makes no difference between v_1 and v_3). It suffices to show that either there is a $v_i \in D$ giving 1 to f or there are two $v_i \in D$'s taking 0 each from f : then the total donation of f is in fact at most $\frac{8}{3} - \frac{2}{3} = 2$. If $v_6 \notin D$, then f receives 1 from v_1 by R4, and we are done. Suppose $v_6 \in D$. Then v_1 , being adjacent to $v_2 \notin D$ and thus having degree at least three, gets nothing from f . Another vertex getting nothing from f is the first among v_6, v_5, \dots that has a neighbor inside D , and we are done again.

Now suppose that neither v_1 nor v_3 belongs to D . Then $d(t) \geq 4$ by (9a). By (9b), either $d(v_i) \geq 4$ or $t \in D$. In the first case, f gives $\frac{1}{6}$ to tv_1v_2 by R0 and nothing to v_1 . It follows that $ch^*(f) \geq 2 - \frac{1}{6} - \frac{1}{2} - \frac{1}{3} \times 4 = 0$. In the second case, f receives 1 across the edge v_1v_2 due to the second part of R4.

This completes the proof for $r(f) = 6$.

The next case is $r(f) = 7$. Now f still is adjacent to at most one 3-face, and so can give at most $\frac{2}{3}$ to a 3-face and at most $\frac{1}{2} \times 2 + \frac{1}{3} \times 5$ to the vertices, i.e., $\frac{10}{3}$ in total. However, giving $\frac{2}{3}$ across an edge implies giving nothing to the two incident vertices of degree at least 4, so in fact $ch^*(f) \geq 0$.

Now suppose $f = v_1 \dots v_r$, where $r \geq 8$. We want to partition all the donation of f to the vertices by R1, R2 and across the edges by R0 into r groups, associated with the incident vertices of f , so that the total donation per group were at most $\frac{1}{2}$. Along with the donation of $(r-8)/2$ to f_0 by R5, this will yield $ch^*(f) \geq r - 4 - r/2 - (r-8)/2 = 0$.

Formally, we define these groups as follows. If f gives $\frac{2}{3}$ across the edge $v_i v_{i+1}$ (addition is mod r), we split this donation of $\frac{2}{3}$ evenly between the vertices v_i and v_{i+1} , which in this case do not receive anything from f . If f gives $\frac{1}{3}$ (or $\frac{1}{6}$) across the edge $v_i v_{i+1}$ by R0, we split this $\frac{1}{3}$ evenly between v_{i-1} and v_{i+2} , which receive at most $\frac{1}{3}$ each

from f by R1, R2 (or, respectively, send $\frac{1}{6}$ to v_{i-1}). It remains to observe that every two crossed edges lie at distance at least 4 along the boundary of f .

$$(12) \text{ } ch^*(f_0) > 0.$$

Recall that $D = d_1 \dots d_7$ and $ch(f_0) = r(f_0) + 4 = 11$. We show that f_0 either sends away less than 11 in total or loses precisely 11 but acquires at least $\frac{3}{2}$ by R5.

By R2, f_0 gives $\frac{5}{3}$ to each incident 2-vertex. By R3, f_0 gives 1 to each incident ≥ 3 -vertex. Since $G \neq C_7$ and G is 2-connected, it follows that D has at least two ≥ 3 -vertices. The total donation of f_0 is thus at most $\frac{2}{3} + 1 \times 2 + \frac{5}{3} \times 5 = 11$, i.e., $ch^*(f_0) \geq 0$. Moreover, the equality can hold only if D is adjacent to a 3-face and incident with five 2-vertices. But then the only non-triangular face f adjacent to f_0 has at least four internal vertices due to the absence of 5-cycles and intersecting 3-faces in G . Thus $r(f) \geq 7 + 4 = 11$, so that f sends to f_0 at least $(11 - 8)/2 > 0$ by R5.

Due to the remark before formulating rules R0–R5, this completes the proof of Theorem 1.4.

Acknowledgments

The authors are indebted to the referees and to A.N. Glebov for carefully reading the manuscript and numerous useful remarks.

References

- [1] M.R. Garey, D.S. Johnson, L.J. Stockmeyer, Some simplified NP-complete graph problems, Theoret. Comput. Sci. 1 (1976) 237–267.
- [2] H. Grötzsch, Ein Dreifarbensatz für dreikreisfreie Netze auf der Kugel, Wiss. Z. Martin Luther Univ. Halle Wittenberg, Math.-Nat. Reihe 8 (1959) 109–120.
- [3] I. Havel, On a conjecture of B. Grünbaum, J. Combin. Theory 7 (1969) 184–186.
- [4] T.R. Jensen, B. Toft, Graph Coloring Problems, Wiley-Interscience, New York, 1995.
- [5] R. Steinberg, The state of the three color problem, in: J. Gimbel, J.W. Kennedy, L.V. Quintas (Eds.), Quo Vadis, Graph Theory? Ann. Discrete Math. 55 (1993) 211–248.